

Supplementary Table 3. Laboratory characteristics of norovirus GII-associated CwG versus enteric adenovirus-associated CwG

Variable	Norovirus GII (N=23)	Enteric adenovirus (N=8)	Total CwG (N=41)	P value
WBC ($10^3/\text{mm}^3$)	9,100.0 (6,300.0–10,800.0)	10,850.0 (8,875.0–15,350.0)	10,000.0 (6,950.0–11,450.0)	0.121
Hemoglobin (g/dL)	11.8 (11.4–12.8)	12.55 (11.93–12.88)	12.1 (11.6–12.8)	0.172
Platelets ($10^3/\text{mm}^3$)	274.0 (243.0–365.0)	310.0 (287.3–457.0)	314.0 (257.0–406.5)	0.172
Sodium (mEq/L)	136.0 (134.0–138.0)	134.5 (134.0–136.8)	136.0 (134.0–138.0)	0.362
Glucose (mg/dL)	81.0 (70.5–90.0)	78.0 (68.0–86.0)	82.5 (71.0–93.8)	0.420
Calcium (mg/dL)	9.7 (9.3–10.2)	9.9 (9.3–10.3)	9.9 (9.5–10.3)	0.842
Uric acid (mg/dL)	8.9 (7.2–9.7)	7.6 (6.8–9.4)	8.1 (5.7–9.7)	0.339
Lactate (mmol/L)	1.6 (1.1–2.3)	2.3 (1.9–3.3)	2.0 (1.3–2.8)	0.030

Values are presented median (interquartile range).

GII, genogroup II; CwG, benign convulsions with mild gastroenteritis; WBC, white blood cell count.

P values indicate comparisons between norovirus GII-associated and enteric adenovirus-associated CwG.

Boldface indicates a statistically significant difference with $P < 0.05$.