

**Supplementary Table 3. Statistical analysis of the three biomarkers (ADM, IL-6, and CRP) at the three collection points (D0, D3, and D7) in the group of 15 nonsurvivors (deaths)**

Parameter	Nonsurvivors (N=15)								
	ADM			IL-6			CRP		
	D0	D3	D7	D0	D3	D7	D0	D3	D7
Minimum	0.46	0.43	0.41	91.86	10.26	60.80	1.90	4.14	3.30
25% Percentile	0.55	0.51	0.47	393.40	139.30	104.14	3.10	7.18	7.12
Median	0.58	0.68	0.61	569.71	302.64	171.60	12.90	17.70	15.05
75% Percentile	0.79	1.22	1.10	732.02	473.20	434.60	18.30	27.38	27.88
Maximum	10.38	11.11	8.73	1,265.0	703.24	796.32	70.12	77.92	79.94
Mean	1.45	2.23	1.715	579.59	338.86	273.00	15.13	21.34	21.30
SD	2.56	3.39	2.49	294.45	217.75	227.74	17.04	20.35	21.76
SEM	0.66	0.88	0.64	76.03	56.22	58.80	4.40	5.26	5.62
Lower 95% CI	0.03	0.35	0.33	416.53	218.28	146.88	5.69	10.07	9.25
Upper 95% CI	2.87	4.11	3.09	742.64	459.45	399.11	24.56	32.61	33.35
CV (%)	176.68	152.19	145.15	50.80	64.26	83.42	112.65	95.37	102.15

ADM, adrenomedullin; IL-6, interleukin 6; CRP, C-reactive protein; LOS, late-onset neonatal sepsis; D0, neonatal intensive care unit admission; D3 and D7, days 3 and 7 after hospitalization and prescription of antibiotics; SD, standard deviation; SEM, standard error of mean; CI, confidence interval; CV, coefficient of variation.